

OVERDOSE ACTION PLAN



OPIOD OVERDOSE ACTION PLAN

How to respond to someone who has overdosed on heroin or other opioids

SIGNS OF OVERDOSE

Not responding
Shallow or no breathing
Blue lips or fingernails
Snoring or gurgling

YES

to any of the signs above proceed to the Action Plan

Whilst care has been taken in preparing the content of this material, the Pharmacy Guild of Australia, its employees and related parties cannot accept any liability—including for any loss, damage or injury—resulting from reliance on the content, or the accuracy, currency or completeness of the material. Produced by Angelo Piccolo and Luke Mitchell

CHECK FOR DANGER

Needles, sharp objects, bystanders etc.
Make sure the environment is safe

CHECK FOR RESPONSE

Call the person's name. Shake their shoulders vigorously or rub their chest with your knuckles



000

If there is no response, or poor response,

SEND FOR HELP


Dial 000 and ask for an ambulance



CLEAR THE PERSON'S AIRWAY

Check their breathing

Watch for their chest rising
Look, listen and feel for 2 breaths in 10 seconds.



IF THEY ARE NOT BREATHING
or their breathing is slow or abnormal

Put them on their back, clear and open their airway, pinch their nose and give two rescue breaths




IF THEY ARE BREATHING NORMALLY


Put them in the recovery position and monitor them until help arrives.
Ensure that their airway is clear and open

**IF THEY ARE STILL NOT BREATHING NORMALLY
GIVE NALOXONE.**

(SEE OVERLEAF FOR PREPARING INJECTION)



Inject the entire contents into muscle of the upper arm or thigh and even through clothing in an emergency



L Note the time given,

Continue rescue breathing—1 breath every 5 seconds

**NALOXONE
WON'T
HARM
A PERSON,
AND IT MAY
SAVE
THEIR
LIFE**

IF THERE IS NO RESPONSE
Inject another dose of Naloxone in 2–5 minutes.

L Note the time given, continue rescue breathing

If you are worried give another dose at any time.

If the person is breathing normally, then put them in the recovery position and monitor them. Ensure that their airway is clear and open.


**If still no response, commence CPR
30 compressions, then 2 breaths; repeat**

For more information, including a list of organisations that provide training visit: www.creidu.edu.au/naloxone

OVERDOSE ACTION PLAN

NALOXONE MINIJET PREFILLED SYRINGE

Remove Naloxone mini-jet from package, unscrew the cap and screw on the 23g needle provided.



IT IS RECOMMENDED THAT YOU HAVE AT LEAST TWO MINISETS ON HAND AS MULTIPLE DOSES MAY BE REQUIRED

NALOXONE AMPOULE

Prepare needle and syringe for muscular injection by unwrapping 3ml syringe and screwing on 23g needle.



USING AN AMPOULE




Take ampoule and hold ampoule between thumb and index finger in one hand



With the other hand, pinch top of ampoule with index finger on blue dot and thumb on other side



Snap ampoule open by breaking the top of the ampoule away from the blue dot



Draw all of the naloxone out of the ampoule by inserting the needle in the naloxone solution and drawing the plunger from the syringe barrel.



IT IS RECOMMENDED THAT YOU HAVE AT LEAST 5 AMPOULES (ONE PACK) ON HAND AS MULTIPLE DOSES MAY BE REQUIRED, AND IT IS POSSIBLE TO BREAK AN AMPOULE ON OPENING